

Plenary paper

Methemoglobinemia and ascorbate deficiency in hemoglobin E β thalassemia: metabolic and clinical implications

Angela Allen,^{1,2} Christopher Fisher,¹ Anuja Premawardhena,³ Dayananda Bandara,⁴ Ashok Perera,⁴ Stephen Allen,² Timothy St Pierre,⁵ Nancy Olivieri,⁶ and David Weatherall¹

¹MRC Molecular Haematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom;

²College of Medicine, Swansea University, Swansea, United Kingdom; ³University of Kelaniya, Colombo, Sri Lanka; ⁴National Thalassaemia Centre, District Hospital, Kurunegala, Sri Lanka; ⁵School of Physics, University of Western Australia, Crawley, Australia; and ⁶Hemoglobinopathy Research, University Health Network, Toronto, ON

During investigations of the phenotypic diversity of hemoglobin (Hb) E β thalassemia, a patient was encountered with persistently high levels of methemoglobin associated with a left-shift in the oxygen dissociation curve, profound ascorbate deficiency, and clinical features of scurvy; these abnormalities were corrected by treatment with vitamin C. Studies of erythropoietin production before and after treatment suggested that, as in an ascorbate-deficient murine model, the hu-

man hypoxia induction factor pathway is not totally dependent on ascorbate levels. A follow-up study of 45 patients with HbE β thalassemia showed that methemoglobin levels were significantly increased and that there was also a significant reduction in plasma ascorbate levels. Haptoglobin levels were significantly reduced, and the high frequency of the 2.2 haptoglobin genotype may place an additional pressure on ascorbate as a free-radical scavenger in this population.

There was, in addition, a highly significant correlation between methemoglobin levels, splenectomy, and factors that modify the degree of globin-chain imbalance. Because methemoglobin levels are modified by several mechanisms and may play a role in both adaptation to anemia and vascular damage, there is a strong case for its further study in other forms of thalassemia and sickle-cell anemia, particularly when splenic function is defective. (*Blood*. 2012;120(15):2939-2944)

Introduction

Although low ascorbate levels have been observed in patients with different hemoglobinopathies,^{1,2} there are very few reports of the clinical manifestations of scurvy in these conditions.³ During an analysis of the mechanisms for the broad phenotypic diversity of hemoglobin E (HbE) β thalassemia in Sri Lanka, a patient was encountered with profound ascorbate deficiency and clinical features of scurvy who also had a high level of methemoglobin. This unusual combination of findings has raised several important questions. First, to what extent does ascorbate deficiency interfere with the hypoxia-sensing mechanism in humans, particularly with respect to erythropoietin response to anemia? The key players in this pathway are the prolyl hydroxylase domain-containing enzymes that catalyze the prolyl-4-hydroxylation of the hypoxia-inducible factor in the presence of oxygen and 2-oxoglutarate as cosubstrates with iron and ascorbic acid as cofactors.⁴⁻⁶ Recent work in ascorbate-deficient mice suggests that other fail-safe mechanisms are involved in this reaction and that erythropoietin response is not altered in ascorbate deficiency^{7,8}; is this the case in humans? The second question raised by this unusual patient report is, because of the potential deleterious effects of methemoglobin on a patient's response to anemia⁹ and the vascular endothelium,¹⁰ (1) how common are increased levels of methemoglobin in HbE β thalassemia and related disorders, (2) to what extent might this depend on ascorbate deficiency, and (3) what other factors may be involved?

The results of these studies suggest that, with respect to hypoxia recognition, humans are able to compensate for ascorbate deficiency in the same way as the murine model. There is a highly significant increase in methemoglobin production in HbE β thalassemia and a significant reduction in plasma ascorbate levels, although not to those observed in the patient whose findings initiated this study. There is, however, a highly significant relationship between the level of methemoglobin and splenectomy and also with the factors that modify globin-chain imbalance. A further complication in this population was the finding that the haptoglobin genotype was nearly all of the 2.2 variety, which is less effective at hemoglobin binding^{11,12} and may place an additional pressure on ascorbate as a free radical scavenger.

Clearly, there are multiple factors involved in the increased level of methemoglobin production in this form of thalassemia and because of its potential effects on adaptation to anemia and vascular damage, further studies of the mechanisms involved in its increased production are required in other types of thalassemia and sickle-cell anemia.

Methods

Patients

The subject in whom the findings initiated this study was a 19-year-old female patient attending the National Thalassaemia Center, Kurunegala, Sri Lanka. She had presented at 2 years of age with anemia and splenomegaly, and later the diagnosis of HbE β thalassemia was established. For the next

Submitted June 12, 2012; accepted July 26, 2012. Prepublished online as *Blood* First Edition paper, August 10, 2012; DOI 10.1182/blood-2012-06-435875.

The publication costs of this article were defrayed in part by page charge

payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2012 by The American Society of Hematology

Table 1. Hematologic and related studies in the propositus, before treatment with vitamin C, and her relatives

Subject	Variable (units), normal range											
	Hb (g/dL), 12-16	MCV (fL), 80-94	MCH (pg), 27-32	HbF (%), < 1	HbA ₂ (%), 2.3-3.2	HbE (%), 0	Met Hb (%), 0.1-0.6	P ₅₀ (mm Hg), 24.1-27.2	Cytb ₅ R (U/gHb), 11.5-26.9	GAPD (IU/gHb), 142.2-308.9	Glutathione reductase (U/gHb), 4.7-13.2	G6PD deficiency screening test
Propositus	6.4	74.3	21.0	18.7		76.7	10.7-13.6	21.3-23.6	50.7	301.5	9.81	Normal
Mother	10.7	54.9	15.8	< 1	4.2		1.9	26.5	28.9	271.0	13.0	Normal
Father	14.2	78.0	23.9	< 1		24.9	0.7	25.2	17.7	230.0	5.5	Normal
Brother	14.6	81.5	28.5	< 1	2.6		0.5	25.2	14.6			Normal
Sister	13.0	84.5	29.6	< 1	2.6		0.4	25.3	21.8			Normal

Cytb₅R indicates cytochrome b₅ reductase; GAPD, glyceraldehyde phosphate dehydrogenase; G6PD, glucose-6-phosphate dehydrogenase; Hb, hemoglobin; MCH, mean cell hemoglobin; Met, methemoglobin; and MCV, mean cell volume.

7 years, she received intermittent transfusion, after which she underwent splenectomy on the basis of a steady-state hemoglobin level of 5-6 g/dL and a spleen that was enlarged to 19 cm below the costal margin. For the next 10 years, despite hemoglobin values in the 7-8 g/dL range, her growth and sexual maturation were delayed, although by the age of 19 years, she had achieved midparental height and the menarche. Only when she reached the age of 17 years did the patient and her family disclose that she had had painful and enlarged gums and intermittent mucosal bleeding for several years. A detailed dietetic history at this time revealed that she ate no fruit of any form or vegetables, a diet that had persisted for several years. The results of a screening for environmental factors that might induce methemoglobinemia were negative.

On the basis of subsequent findings in this patient and her family, detailed studies were conducted on 45 patients attending the National Thalassemia Center who were chosen at random from more than 200 patients with HbE β thalassemia who were being followed at the Center. Clinical and hematologic data on this group of patients have been published previously, together with a detailed account of a classification system directed at defining the phenotypic variability of the disease.¹³ In short, a "mild" phenotype covers those who never required transfusion, had stopped transfusion with no ill effects after several years follow-up, or had required no further transfusion after splenectomy. Patients with "severe" phenotypes were defined as those who relied on long-term transfusion. There were 25 of the former and 20 of the latter in this study. In addition, methemoglobin values were estimated in 17 normal adult volunteers and 17 patients with different types of hemoglobinopathy at the Center.

Procedures

Venous blood was collected into heparin and EDTA from all study participants. Duplicate measurements of methemoglobin levels and P₅₀ were made from the heparinized blood sample with the use of a Rapidpoint 405 analyzer with an integral co-oximeter (Bayer). This instrument incorporates a polychromator that allows the simultaneous measurements across the various fractions of hemoglobin in the range of 473-671 nm. To confirm that methemoglobin levels were being measured accurately, blood samples from the propositus and a group of patients with HbE β thalassemia and normal controls were analyzed via the manual method of Evelyn and Malloy.¹⁴ We found there was close agreement between the methemoglobin values between the 2 methods used.

The samples were then centrifuged, the plasma removed, and plasma ascorbate levels measured immediately with a ferric-reducing ascorbate assay (procedure K671; BioVision). To prevent plasma protein precipitation, the ferric-reducing ascorbate buffer was diluted 1 in 10 before use. Plasma haptoglobin was measured with the use of a commercial assay (procedure TP.801; Tridelata Development Ltd). Routine hematologic indices were measured in the EDTA sample (Coulter Electronics). The sample was centrifuged, the plasma removed from the cells, and both were stored at -20°C until shipped to Oxford on dry ice. Plasma erythropoietin and IL-8 levels were measured with an enzyme-linked immunosorbent assay kit (DEP00; R&D Systems) and a Compact human IL-8 ELISA kit (M1918; PeliKine). DNA was extracted from the cell pellet with the use of a QIAGEN DNA blood mini kit (51104), and the haptoglobin genotype was

determined by polymerase chain reaction.¹⁵ Hemoglobin analysis, serum ferritin levels, and hepatic iron concentrations (measured by magnetic resonance imaging) followed previously reported methods.^{16,17}

To investigate the patient with a markedly increased methemoglobin concentration, further blood samples were collected from her and her immediate family and transferred into EDTA and acid citrate dextrose. EDTA samples were screened for glucose-6-phosphate dehydrogenase (G6PD) deficiency with the use of a qualitative assay (procedure 400; Trinity Biotech). Levels of reduced glutathione were determined (kit 371757; Calbiochem) and a red cell hemolysate, stabilized in EDTA-mercaptoethanol, was prepared and used for the measurement of cytochrome b₅ reductase and glyceraldehyde phosphate dehydrogenase.¹⁸ A red cell hemolysate was prepared from each acid citrate dextrose sample, and G6PD activity was measured with a quantitative ultraviolet, kinetic assay (procedure 345-uv; Trinity Biotech Co). Glutathione reductase was measured using a quantitative manual method (kit GR2368; Randox Laboratories). Pyruvate kinase was measured according to the method described by Dacie and Lewis.¹⁹ Both cytochrome b₅ and cytochrome b₅ reductase genes and the *HBA* and *HBB* genes were sequenced.²⁰

A urine sample was collected from the patient and tested with Combur 10 diagnostic strips (Roche Diagnostics) for the presence of nitrites and hemoglobin. The urine sediment was examined by microscopy for the presence of red blood cells.

Statistical analysis

Statistical analysis was performed with SPSS 16.0 for Windows (Release 16.0.1; SPSS Inc). Differences between median methemoglobin concentrations were assessed with the Mann-Whitney *U* test. We used multiple regression analysis to explore the relationship between methemoglobin and the potential predictor variables severity, transfusion status, and splenectomy. *P* < .05 was considered statistically significant.

Ethical approval

Approval for the research program on HbE β thalassemia was obtained from the Ethical Committee of the College of Pediatricians, Colombo, Sri Lanka, and the Oxford Tropical Research Ethical Committee, Oxford, United Kingdom. This study was conducted in accordance with the Declaration of Helsinki.

Results

The findings in the family that led to these studies are summarized in Table 1 and further biochemical data of the propositus, including findings before and after treatment, in Table 2. The propositus had a hemoglobin pattern typical of HbE β thalassemia; sequencing of the *HBB* genes revealed the β^E mutation on 1 chromosome and the severe β thalassemia mutation, IVS1-5 (G-C), which is very common in the Sri Lankan population,¹⁶ on the other. Further

Table 2. Further biochemical and related analyses of the propositus, including, in some cases, data obtained before and after treatment with vitamin C

Treatment period	Variable (units), normal range											
	Ascorbate (nmol/mL), 28-84	Met Hb (%), 0.1-0.6	P ₅₀ (mm Hg), 24.1-27.2	Mean Hb (g/dL), 12-16	Mean Epo (IU/mL), 2-14	G6PD (U/gHb), 4.6-13.5	Pyruvate kinase (IU/gHb), 7.2-14.0	GSH, (μmol/gHb), 6.6-10.0	Haptoglobin (g/dL), 0.3-2.0	Hepatic iron (mg/g dw), 0.6-1.2	IL-8, (pg/mL) < 10	Urinary nitrite screen
Pretreatment	2.1-6.7	10.7-13.6	21.3-23.6	6.2	94.3	21.1	7.7	6.18	0.26	2.9	179.6	Negative
Posttreatment	30.2-36.6	0.85	26.9	7.0	76.8							

Hb and Epo values are the means of 5 estimations.

Epo indicates erythropoietin; G6PD, glucose-6-phosphate dehydrogenase; Hb, hemoglobin; and Met, methemoglobin.

sequencing of the *HBA* and *HBB* genes revealed no other abnormalities, excluding hemoglobin M mutations. The mother had findings typical of β thalassemia trait, although complicated by iron deficiency anemia because of menorrhagia. The father had hemoglobin E trait, whereas the patient's 2 siblings were healthy.

Multiple estimations indicated that the propositus had markedly increased levels of methemoglobin in the range of 10.7%-13.6%. The patient's mother with β thalassemia trait had a slightly increased level of methemoglobin, whereas the levels in other family members were normal. The cytochrome b₅ reductase levels were increased in the propositus and normal in the other family members. Both cytochrome b₅ (*cytb5*) and cytochrome b₅ reductase (*cytb5r*) genes from the propositus were sequenced and showed no abnormality. Glyceraldehyde phosphate dehydrogenase and glutathione reductase levels were within the normal range in the propositus and both her parents. The patient's P₅₀ was significantly reduced compared with other family members, resulting in a marked left shift in the oxygen-dissociation curve compared with those of patients in the same population with HbE β thalassemia recently reported.²¹ Screening for glucose-6-phosphate dehydrogenase deficiency was negative in all family members.

To determine the reason for the extremely high methemoglobin level in the propositus, we performed further investigations, as shown in Table 2. Her plasma ascorbate level was extremely low, whereas G6PD levels were increased. Reduced glutathione levels were at the lower limit of the normal range. Haptoglobin levels were slightly reduced, and there was an increased level of IL-8. Treatment with vitamin C, 50 mg on alternate days, resulted in a dramatic decrease in the methemoglobin level and an increase in

the ascorbate level to above the lower range of normal. There was also a shift to the right in the oxygen dissociation curve associated with a significant increase in the P₅₀. There was no significant change in the relationship between hemoglobin and erythropoietin levels estimated on 5 samples before and after treatment.

To explore further the significance of the findings in this patient, 45 patients with HbE β thalassemia were chosen at random from the 200 or more patients with this condition who were being followed in Kurenegala. Their division into strictly defined severity groups, as described in "Procedures," main clinical and hematologic findings and some of the genetic modifiers responsible for the variation in their phenotypic severity have been reported previously.^{13,16} The major findings in these patients in relationship to the present study are summarized in Tables 3 and 4. As shown in Table 3, those with HbE β thalassemia had a significant increase in methemoglobin compared with healthy controls and univariate analysis showed that there was a highly significant increase in methemoglobin levels in those who had been splenectomized compared with those who had intact spleens. As shown in Table 3, there was also a significant relationship between methemoglobin levels and phenotypic severity, as judged by the findings in the mild and severe groups and mirrored by the transfusion requirements. In multiple regression analysis, only splenectomy was statistically significantly related to methemoglobin level (standardized $\beta = 0.64$, $t = 3.68$, $P < .01$).

As shown in Table 4, the mean plasma ascorbate level in this group of patients was at the bottom limit of the normal range; in 10 cases it was subnormal. Although the number of cases available with matched plasma ascorbate and methemoglobin levels was too

Table 3. Analysis of methemoglobin levels in 45 patients with HbE β thalassemia with a breakdown of cases into splenectomized and nonsplenectomized, low and high transfusion rates, and mild and severe phenotypes as defined in the text

Diagnosis	n	Median Met Hb, %	Interquartile range	Range	P
Normal controls	17	0.3	0.25-0.4	0.1-0.6	< .001
HbE β thalassaemia (all cases)	45	2.7	1.9-3.65	0.9-6.3	
Hb E β thalassaemia (splenectomized)	20	3.7	3.1-4.2	0.9-6.3	< .001
HbE β thalassaemia (spleen intact)	25	2.3	1.5-2.8	0.9-4.8	
Hb E β thalassaemia (mild)	25	2.45	1.9-3.6	0.9-4.8	.084
HbE β thalassaemia (severe)	20	3.1	1.9-3.9	0.9-6.3	
Hb E β thalassaemia (0-20 blood transfusions)	21	2.5	1.85-3.1	0.9-3.6	.001
HbE β thalassaemia (> 20 blood transfusions)	18	3.65	3.4-4.5	1.6-6.3	
HbE β thalassaemia (mild, spleen intact)	13	2.2	1.75-2.5	0.9-3.6	.018
HbE β thalassaemia (mild, splenectomized)	12	3.6	2.65-4.05	0.9-5.0	
Hb E β thalassaemia (severe, spleen intact)	12	2.65	1.38-3.18	0.9-4.8	.002
HbE β thalassaemia (severe, splenectomized)	8	4.0	3.7-4.45	3.5-6.3	
Hb E β thalassaemia (mild, spleen intact)	13	2.2	1.75-2.5	0.9-3.6	.32
HbE β thalassaemia (severe, spleen intact)	12	2.65	1.38-3.18	0.9-4.8	
Hb E β thalassaemia (mild, splenectomized)	12	3.6	2.65-4.05	0.9-5.0	.01
HbE β thalassaemia (severe, splenectomized)	8	4.0	3.7-4.45	3.5-6.3	

Hb, hemoglobin; and Met, methemoglobin.

Table 4. Additional data from the 45 patients with HbE β thalassemia, including ascorbate and iron status and levels of IL-8 and haptoglobin, including the different genetic forms of the latter

Variable (units), normal range	n	Median	Interquartile range	Range
Methemoglobin (%), 0.1-0.6	45	2.70	1.9-3.65	0.9-6.3
Plasma ascorbate (nmol/mL), 28-84	20	28.23	21.03-31.68	12.6-56.3
IL-8 (pg/mL), < 10	45	7.97	3.47-172.8	0.39-2210
Hepatic iron (mg/g dw), 0.6-1.2	33	6.0	2.95-10.75	1.0-33.0
Haptoglobin (g/L), 0.3-2.0				
All cases	38	0.24	0.20-0.28	0.16-0.42
Haptoglobin genotype 1:1	0			
Haptoglobin genotype 2:1	8	0.27	0.26-0.33	0.22-0.33
Haptoglobin genotype 2:2	25	0.23	0.19-0.26	0.16-0.41

Hb indicates hemoglobin.

small to determine whether there was a significant relationship between the 2 at this level of plasma ascorbate, the mean level of methemoglobin in those in whom these measurements were available and who had all been splenectomized was 4.24%, and the mean level of plasma ascorbate was 23.1 nmol/mL, that is in the subnormal range.

Overall, the level of haptoglobin was subnormal in the patients with HbE β thalassemia, although no case of absent haptoglobin was encountered. The majority of the patients had the 2.2 haptoglobin genotype. There was a wide range of hepatic iron concentrations that were not significantly related to the methemoglobin level. There was also a highly significant elevation of IL-8 levels in this patient population; levels were elevated (> 10 pg/mL) in 21 cases, with some having very high levels (Table 4).

The findings in the other hemoglobin disorders studied (Table 5) suggest further studies of patients with β thalassemia major or intermedia and sickle cell anemia and related conditions are indicated. The rare sickle cell disorders in Sri Lanka all have the Asian haplotype, which is associated with a relatively mild phenotype.

Discussion

There seems little doubt that the high level of methemoglobin in the patient whose findings initiated these studies resulted from profound ascorbate deficiency. It has been estimated that ascorbate is responsible for approximately 16% of methemoglobin reduction in red cells,²² with the remainder relying on several enzymes, notably

Table 5. Miscellaneous methemoglobin levels in conditions related to HbE β thalassemia obtained from patients attending the same clinic

Diagnosis	No.	Methemoglobin, %
Normal, median; range	17	0.3; 0.1-0.6
β thalassemia intermedia	4	1.4, 2.9, 4.7, 4.8
Hb E/ $\delta\beta$ thalassemia	1	2.4
$\delta\beta/\beta$ thalassemia	3	2.0, 2.8, 2.8
Hb SS disease	2	0.3, 1.2
Hb S/ β thalassemia	1	1.8
Hb SD disease	1	1.2
Hb SE disease	1	1.4
β thalassemia trait	2	0.4, 0.5
Hb E trait	2	0.3, 0.3

Hb indicates hemoglobin.

cytochrome b₅ reductase, glyceraldehyde-3-phosphate dehydrogenase, and glutathione reductase.²³ The levels of these enzymes were all increased or normal in the propositus and her family members, and structural studies of the cytochrome b₅ reductase gene of the propositus were normal. Sequencing of the *HBA* and *HBB* genes excluded HbM. Furthermore, the administration of cautious doses of ascorbate, because of the possible deleterious effects of rapid iron mobilization,²⁴ rapidly reversed the methemoglobin levels into the low-normal range. Before treatment the propositus also had a low P₅₀ and a marked left shift in her oxygen dissociation curve, a finding that has been previously observed in association with increased methemoglobin levels.⁹ The ferric (Fe³⁺) hemes are unable to reversibly bind oxygen, and they increase the oxygen affinity of the associated ferrous hemes (Fe²⁺) in the hemoglobin tetramer, causing a left shift in the oxygen dissociation curve.²⁵ There was a significant increase in the P₅₀ with a right shift in the oxygen dissociation curve after treatment with ascorbate. In short, this reflects a more effective adaptation to anemia, as recently described in patients with HbE β thalassemia.²¹

The finding that multiple estimations of the erythropoietin response to a particular hemoglobin level did not change in the propositus before and after treatment with ascorbate is of particular interest with regard to response to hypoxia. These findings are similar to those reported recently in an ascorbate-deficient mouse model.^{7,8} An increased erythropoietin response to anemia depends on the oxygen-sensing properties of the prolylhydroxylase domain-containing enzymes that catalyze the prolyl-4-hydroxylation of the hypoxia-inducible factor and require oxygen and 2-oxoglutarate as cosubstrates with iron and ascorbic acid as cofactors.^{4,6} The erythropoietin response to anemia in the murine model was normal despite profound ascorbate deficiency.⁸ A major compensatory mechanism appeared to be the action of reduced glutathione, the levels of which remained normal or increased in the propositus in the present study.⁸ These observations suggest that the relationship between ascorbate and hypoxia response in humans and mice are similar.

What are the broader issues resulting from these findings? In particular, because the results of the studies in this unusual patient provide clear evidence that ascorbate deficiency can induce methemoglobinemia in HbE β thalassemia, how common are increased methemoglobin levels in this condition and are the levels related mainly to ascorbate or are other factors involved? There have been relatively few reports of the levels of methemoglobin in the inherited hemoglobin disorders. An early study of a few cases of HbE β thalassemia in northern India suggested that methemoglobin levels might be increased in this condition,²⁶ and increased levels have been reported in some cases of inherited unstable hemoglobins²⁷ and sickle cell anemia.^{28,29}

In the present study, there was a significant increase of methemoglobin in a group of patients with HbE β thalassemia whose mean level of plasma ascorbate was at the lower limit of normal; 10 cases showed subnormal levels. However, no cases were encountered with a reduction to the level found in the propositus in this study and the extent to which ascorbate deficiency may be responsible for the modest increase in methemoglobin in these patients requires further study. The most striking finding, however, was the highly significant relationship between splenectomy and methemoglobin levels together with the effect of phenotypic severity, including blood transfusion status. Because the main factors underlying phenotypic variability in this group of patients identified so far are the coinheritance of α thalassemia or relatively high levels of HbF,^{13,16,30} both of which modify the

degree of globin-chain imbalance, it seems likely that splenic function and the degree of excess α -chain synthesis play a major role in determining the level of methemoglobin, at least in HbE β thalassemia.

What is the source of the increased methemoglobin? As in other forms of β thalassemia, excess α -chains are produced in HbE thalassemia with the production of red cell inclusions³¹; despite the mild instability of HbE β^E chains are not found in these precipitates.³² One of the major degradation products of excess α -chains are hemichromes, which bind to the red cell membrane and promote sequestering of band 3.³³ As they form, they go through reversible and irreversible phases during which methemoglobin is produced as an intermediate. It is possible, therefore, that abnormal red cells exposed to this mechanism are recognized and sequestered in the spleen and hence the level of methemoglobin is increased after splenectomy. Because, like other forms of thalassemia, there is a significant hemolytic component in HbE β thalassemia, it follows that the circulation will be continuously exposed to increased levels of methemoglobin.

Another potential source of methemoglobin, in this case in plasma, is the further oxidation of hemoglobin released during hemolysis, the fail-safe mechanism in this case again is binding by haptoglobin. In the present study the haptoglobin levels were reduced in the patients with HbE β thalassemia, although only to a minor degree. However, molecular analysis showed that in almost every case the haptoglobins were of the 2.2 variety, which has been shown to be less effective than the 1.1 variety with respect to hemoglobin binding and which occurs commonly in some Asian countries.³⁴ Recent studies suggest that because of its reduced binding properties, it may put greater pressure on the use of ascorbate as a free radical scavenger and, indeed, may be associated with increased frequency of the clinical manifestations of ascorbate deficiency.^{11,12,35}

Methemoglobin is a significant activator of endothelial cells by stimulation of E-selectin, IL-6, and IL-8 production.¹⁰ It is of

interest therefore that the IL-8 levels in this series of patients with HbE β thalassemia were considerably increased. Because of increasing evidence for vascular complications in other forms of thalassemia intermedia³⁶ and in sickle cell disease, and because of the results of the small pilot study shown in Table 5, further investigation of the potential pathologic role of methemoglobin is indicated, particularly in conditions with reduced splenic function or in which splenectomy is commonly practiced.

Acknowledgments

The authors thank Melanie Percy and Terry Lappin for their advice and Jeanne Packer and Liz Rose for their help in preparing the manuscript.

This work was supported by grants from the United Kingdom Wellcome Trust and Medical Research Council, the U S March of Dimes, and The Anthony Cerami and Ann Dunne Foundation for World Health.

Authorship

Contribution: A.A. and C.F. performed the laboratory studies; S.A. conducted the statistical analysis; A. Premawardhena, D.B., A. Perera, T.S.P., and N.O. collected and analyzed the clinical data on the patients, and A.A. and D.J.W. designed the study and wrote the manuscript.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Professor Sir David Weatherall, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Headington, Oxford, OX3 9DS, United Kingdom; e-mail: liz.rose@imm.ox.ac.uk.

References

- Chiu D, Vichinsky E, Ho SL, Liu T, Lubin BH. Vitamin C deficiency in patients with sickle cell anemia. *Am J Pediatr Hematol Oncol*. 1990;12(3):262-267.
- Claster S, Wood JC, Noetzi L, et al. Nutritional deficiencies in iron overloaded patients with hemoglobinopathies. *Am J Hematol*. 2009;84(6):344-348.
- Cohen A, Cohen IJ, Schwartz E. Scurvy and altered iron stores in thalassemia major. *N Engl J Med*. 1981;304:158-160.
- Lee FS, Percy MJ. The HIF pathway and erythrocytosis. *Ann Rev Pathol*. 2011;6:165-192.
- Schofield CJ, Ratcliffe PJ. Oxygen sensing by HIF hydroxylases. *Nat Rev Mol Cell Biol*. 2004;5(5):343-354.
- Webb JD, Coleman ML, Pugh CW. Hypoxia, hypoxia-inducible factors (HIF), HIF hydroxylases and oxygen sensing. *Cell Mol Life Sci*. 2009;66(22):3539-3554.
- Lappin T, Masson N. Two antioxidants are better than one. *Blood*. 2011;117(20):5276-5277.
- Nytko KJ, Maeda N, Schlafli P, Spielmann P, Wenger RH, Stiehl DP. Vitamin C is dispensable for oxygen sensing in vivo. *Blood*. 2011;117(20):5485-5493.
- Darling RC, Roughton FJW. The effect of methemoglobin on the equilibrium between oxygen and hemoglobin. *Am J Physiol*. 1942;137:56-68.
- Liu X, Spolarics Z. Methemoglobin is a potent activator of endothelial cells by stimulating IL-6 and IL-8 production and E-selectin membrane expression. *Am J Physiol Cell Physiol*. 2003;285(5):C1036-C1046.
- Cahill LE, El-Soehy A. Haptoglobin genotype modifies the association between dietary vitamin C and serum ascorbic acid deficiency. *Am J Clin Nutr*. 2010;92(6):1494-1500.
- Delanghe JR, Langlois MR, De Buyzere ML, Torck MA. Vitamin C deficiency and scurvy are not only a dietary problem but are codetermined by the haptoglobin polymorphism. *Clin Chem*. 2007;53(8):1397-1400.
- Premawardhena A, Fisher CA, Olivieri NF, et al. Haemoglobin E beta thalassaemia in Sri Lanka. *Lancet*. 2005;366:1467-1470.
- Dacie JV, Lewis SM. *Practical Haematology*. 7th ed. Edinburgh: Churchill Livingstone; 1991:191.
- Imrie H, Fowkes FJ, Michon P, et al. Haptoglobin levels are associated with haptoglobin genotype and alpha+ -Thalassaemia in a malaria-endemic area. *Am J Trop Med Hyg*. 2006;74(6):965-971.
- Fisher CA, Premawardhena A, de Silva S, et al. The molecular basis for the thalassaemias in Sri Lanka. *Br J Haematol*. 2003;121(4):662-671.
- St. Pierre T, Olivieri N, Thayalasuthan V, et al. Relationship between serum ferritin and liver iron concentration in hemoglobin E thalassemia. *Haematologica*. 2011;96(2):69.
- Beutler E. *Red Cell Metabolism: A Manual of Biochemical Methods*. 3rd ed. New York: Grune & Stratton; 1984:81-82.
- Dacie JV, Lewis SM. *Practical Haematology*. 7th ed. Edinburgh: Churchill Livingstone; 1991:214-215.
- Percy MJ, Gillespie MJ, Savage G, Hughes AE, McMullin MF, Lappin TR. Familial idiopathic methemoglobinemia revisited: original cases reveal 2 novel mutations in NADH-cytochrome b5 reductase. *Blood*. 2002;100(10):3447-3449.
- Allen A, Fisher C, Premawardhena A, et al. Adaptation to anemia in hemoglobin E-beta thalassemia. *Blood*. 2010;116(24):5368-5370.
- Jaffé ER, Neumann G. Hereditary methemoglobinemia, toxic methemoglobinemia and the reduction of methemoglobin. *Ann N Y Acad Sci*. 1968;151(2):795-806.
- Nagel RL, Jaffe ER. CO-, NO-, Met-, and Sulf-hemoglobinemias: the dyshemoglobins. In: Steinberg MH, Forget BG, Higgs DR, Nagel RL, eds. *Disorders of Hemoglobin*. 1st Ed. Cambridge: Cambridge University Press; 2001:1214-1234.
- Nienhuis AW. Vitamin C and iron. *N Engl J Med*. 1981;304:170-171.
- Agarwal N, Nagel RL, Prchal JT. Dyshemoglobinemias. In: Steinberg MH, Forget BG, Higgs DR, Weatherall DJ, eds. *Disorders of Hemoglobin*. 2nd Ed. New York: Cambridge University Press; 2009:607-622.
- Swarup S, Ghosh SK, Chatterjea JB. Methaemoglobin level and its relation to the stability of erythrocytic reduced glutathione in thalassaemia syndrome. *Indian J Med Res*. 1964;52:273-278.
- Dacie JV. *The Unstable Haemoglobin Haemolytic*

- Anaemias. Vol. 2 The Haemolytic Anaemias.* 3rd Ed. London: Churchill Livingstone; 1988:348-349.
28. Caboot JB, Jawad AF, McDonough JM, et al. Non-invasive measurements of carboxyhemoglobin and methemoglobin in children with sickle cell disease. *Pediatric Pulmonol.* 2012;47(8):808-815.
 29. Zerez CR, Lachant NA, Tanaka KR. Impaired erythrocyte methemoglobin reduction in sickle cell disease: dependence of methemoglobin reduction on reduced nicotinamide adenine dinucleotide content. *Blood.* 1990;76(5):1008-1014.
 30. Fucharoen S, Weatherall DJ. Hemoglobin E disorders. In: Steinberg MH, Forget BG, Higgs DR, Weatherall DJ, eds. *Disorders of Hemoglobin.* 2nd Ed. New York: Cambridge University Press; 2009:417-433.
 31. Weatherall DJ, Clegg JB. *The Thalassemia Syndromes.* 4th Ed. Oxford: Blackwell Science; 2001.
 32. Wickramasinghe SN, Lee MJ. Observations on the relationship between gamma-globin chain content and globin chain precipitation in thalassaemic erythroblasts and on the composition of erythroblastic inclusions in Hb E/beta-thalassaemia. *Eur J Haematol.* 1997;59:305-309.
 33. Rachmilewitz EA, Schrier SL. Pathophysiology of beta thalassemia. In: Steinberg MH, Forget BG, Higgs DR, Nagel RL, eds. *Disorders of Hemoglobin.* 1st Ed. Cambridge: Cambridge University Press; 2001:233-251.
 34. Langlois MR, Delanghe JR. Biological and clinical significance of haptoglobin polymorphism in humans. *Clin Chem.* 1996;42(10):1589-1600.
 35. Delanghe JR, Langlois MR, De Buyzere ML, et al. Vitamin C deficiency: more than just a nutritional disorder. *Genes Nutr.* 2011;6(4):341-346.
 36. Taher AT, Musallam KM, Cappellini MD, Weatherall DJ. Optimal management of beta thalassaemia intermedia. *Br J Haematol.* 2011; 152(5):512-523.



blood

2012 120: 2939-2944
doi:10.1182/blood-2012-06-435875 originally published
online August 10, 2012

Methemoglobinemia and ascorbate deficiency in hemoglobin E β thalassemia: metabolic and clinical implications

Angela Allen, Christopher Fisher, Anuja Premawardhena, Dayananda Bandara, Ashok Perera, Stephen Allen, Timothy St Pierre, Nancy Olivieri and David Weatherall

Updated information and services can be found at:
<http://www.bloodjournal.org/content/120/15/2939.full.html>

Articles on similar topics can be found in the following Blood collections

- [Free Research Articles](#) (3039 articles)
- [Plenary Papers](#) (448 articles)
- [Red Cells, Iron, and Erythropoiesis](#) (629 articles)
- [Sickle Cell Disease](#) (99 articles)

Information about reproducing this article in parts or in its entirety may be found online at:
http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:
<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:
<http://www.bloodjournal.org/site/subscriptions/index.xhtml>